# **MERRA**

Update for the User Group

25 August, 2006


## Overview

- Progress since March (last update to User Group)
- Issues we are addressing
- Schedule
- Advice sought from User Group
- Appendix (Validation strategy; Sweeper strategy; Forward processing stream strategy)

# GEOS-5 DAS Progress since March


- 1/2° eros system with IAU running in real-time parallel operations since June
- Non-science data-sweep in progress -- resolving issues with data, CRTM, QC
- 2º system used for 2001 science-quality tests -- addressing issues
  - Physics tuning in DAS model has improved moisture, precipitation and cloud fields
  - Online model bias correction through the DAS has been implemented and initial tests positive
  - Tuned background statistics and implemented balance constraint will improve masswind balance
  - Implemented re-play mode for tuning model and analysis
  - Aerosols added to radiation code
- Technical improvements: single executable; improved I/O performance
- MERRA file spec has additions to operational file spec -- near finalized

### Zonal Mean Temperature for January 2001




#### 1/2° tests

### Zonal Mean Temperature for January 2006




**Problem corrected in operations** 


### July 2001 Precipitation (mm/day)




## July 2001 Precipitation - CMAP (mm/day)




## July 2001 OLR (Wm<sup>-2</sup>)




## July 2001 LW CRF (Wm-2)




## July 2001 TPW (mm)




## July 2001 TPW - NVAP (mm)



#### Precipitation & Online Moisture Bias Correction




#### July 2001: Impact of SSMI - Zonal mean Specific Humidity




Base: No SSMI

GPROF: SSMI rain-rate retrievals SSMI rad: SSMI radiances only

## January 2001: Impact of SSMI





## JAN 2001 SSM/T2 Observed and Simulated 183.3 +/-1 GHz Tbs (K)

- Sensitive to moisture in
  200 500 hPa layer
- High (low) Tbs signify dry (moist) regions.

From Pete Robertson, MSFC

#### **MERRA**

## Requirement to justify proceeding -- advance upon ERA-40

- Improved treatment of changing observing system
  - SSMI jump evident in other reanalyses has to be reduced
  - Strategy: retrospective bias correction through the model for MERRA

#### Improved Hydrological cycle

- ✓ Minor tuning remains but Tropical Precipitation and TPW already an improvement on era-40
- ✓ Improved cloud-radiation interaction and surface/TOA radiative budgets
- Is vertical distribution of moisture good enough?
- Can we use precipitation data more effectively?

#### Surface


Surface temperature has warm bias (currently no surface analysis)

#### √ Improved stratospheric analysis


- ✓ fully resolved middle atmosphere an advance on era-40
- √ fully interactive ozone an advance on era-40
- ✓ stratospheric transport less noisy than era-40

### January 2001 Specific Humidity (g/kg)


Jan. 2001 Specific Humidity - ERA40 (g/kg)




Jan. 2001 Specific Humidity - NVAP (g/kg)



#### January 2001 T2m - CRU (K)



# **Precipitation Trends**



Precipitation

5.5

90S-90N

30S-30N

4.5

2.5

1960 1965 1970 1975 1980 1985 1990 1995 2000

Time (year)

Satellite era Starts Wippe)


ERA-40

Our goal for MERRA: consistent moisture and precipitation throughout the re-analysis period

**Strategy** (based on suggestion from Kevin Trenberth): Correct the model moisture bias throughout the integration

- Methodology: Use online bias estimation from DAS to obtain diurnal and seasonal components of moisture bias correction for a recent multi-year assimilation (use 1 degree resolution).
- Validation: Repeat training period using fixed correction; test pre-SSMI period.

#### **Schedule**



Expected duration of task
Duration of task with unexpected delays (computer outages, other issues)
4 seasons only for validation

# MERRA UPDATE: Issues for User Group

#### Schedule

■ Goal is to start production in April 2007 which would imply completion spring 2009

#### When do we stop developing?

- Performance targets
- Is delay for moisture bias correction worth it?
- Decision points
  - mid-December: assess progress in bias correction
  - early January: 4-month vs 12-month validation decision


# Appendix Background slides

# Validation Strategy

- Pre-Sweeper tests on 3 periods (1998, 2001 and 2004)
   primarily for data
  - > 2001: Re-Pre-Sweeper includes IAU and Eros
- Sweeper Science run, regular comparisons with existing reanalyses (ERA40, NCEP R2, JRA25 and GEOS4)
- Validation Experiments 2001 & 2004
  - ½ degree, includes metrics for instrument team needs, forecast skills
  - 2004: Comparisons with operational analyses available via CEOP model output data archive

# Sweeper Execution

- Purpose: Exercise the data; Provide MERRA IC; Climate First-look
  - Currently a Data Sweeper running
- AMIP Climate simulation provides stream initialization 5 years upstream from the ½ degree start time
- Minimum 10-15 years of execution needed for ½ degree initialization



## MERRA Execution

- One year spin up at ½ degree from sweeper ICs
- Streams begin: Jan 1 1979, 1989 and 1999
- Along side MERRA
  - Reduced Observing System Baseline (ROSB) using conventional and TOVS data (low priority queue)
  - > AMIP run

